Title: "Guava (Psidium guajava) 13-Hydroperoxide Lyáse and Usés Thereof:" Inventor: Whitehead et al. Attorney Docket No. 06027.0001U3




FIG.1A

Title: "Guava (Psidium guajava) 13-Hydroperoxide Lyase and Uses Thereof:" Inventor: Whitehead et al. Attorney Docket No. 06027.0001U3




FIG.1B

Title: "Guava (Psidium guajava) 13-Hydroperoxide Lyase and Uses Thereoft." Inventor: Whitehead et al. Attorney Docket No. 06027.0001U3

## 3/4

	3/4				
252	25 -	윤율노	8 H - 5	<b>≅</b> ≡-	웃음~
<b>₽</b> = _	88 -	7 × 510 ×	- E - E - E - E - E - E - E - E - E - E	249 Z ATT 1 —	88 H
≈ 8 ~	88 DS >-	<b>多</b> 8 9 0	198 199 7 F CCC 1	248 2 GK A E	C 2 2 2 3 3 3 3
7 3 3 5 4 5	5 219	147 PS TS	196 197 198 199 CAA ANG TIC CTC 0 K F L Peptide	C 22 —	297 Z
<u> </u>	% 95 %	- 99	86 86 C	246 247 CTA GTG L V	= 55 38
9 25 _ 9 25 _ 9 26 _	왕당 >	CAN CAC C	85 55 7	310	₩ ₩ ₩
# 25 G	8 골 =	7 5 6 5 7 6 5 7 7 9	P 50 P	3 244 245 246 24 5 CAG CCT CTA G	25 A A A A
7 8 =	250	200	887	200	2337
7 23 3	26.35 ×	142 143 1 100 996 ( S E	120 19 11 19 11 19	7 6 7	262 26
± 35 ≻ 0 0 0 ×		E 25 -	191 192 193 1 CIC ATC TIC C V I F Ptide 12	7 5 5	2 2 3 E
8 8 °	90 91 AVC G/G V	문용으	20 S	0.00	280 Z
885	888	25 - 25 -	189 190 191 192 193 194 6CC ACC CIC ATC TC CCT A S V   F P Peptide 12	2.58 240 241 242 243 24 C inc att gec gir cic co	288 78 1 288 78
~ <b>5</b> ~	882	22 Z	20 M =	238.2 AIC M	25.25
	88 E E E	137 13 A	181 30	237 Z3 10 Z4 Z4 1 Z4	287 28 TIC 00
25 AC 57 - AC 57 - AC 57	887	25 25 20 20 20 20 20 20 20 20 20 20 20 20 20	25 55 C	236 Z P 200 Z P 200 Z	28 28 U
~ Z ~		200	88 50 ×	223	285 28 600 GV
	\$8 ±3 1 1 3 1 3 1 1 1 1 1 1 1 1 1 1 1 1 1 1	200 0	± 3 × =	78 782 50 010 1 1	× 35 32
₹35 <b>~</b> ≈35 ~	# 25 J	25 24 AC 36 - R	133 150 150 150 150 150 150 150 150 150 150	23.23	282
	82 83 001 000 P P	122 132 132 132 132 132 132 133 133 133	182 18 193 CI 15 CI 15 CI	222 23 C1C 23 L 0	25 25 25 25 26 25 27 26
20 10 10 10 10 10 10 10 10 10 10 10 10 10		55 55 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	# 9 °	257	22 22
~ 5 - ~ 5 -	0 81 0 00 0 00 0 00	25 SS -	883g 2012 S	230 231 2 CTC 0CT (	85
200 Z 200 Z 200 Z	83 ×	129 13 35 AC 17 AC	∞ 55 <sup>ω</sup>	229 23 100 C1	279 280 281 300 GTC GAA A V E
	200 - 200 200 -	128 X	E N .	Z28 22 200 10 R 10	225
22 Z 22 Z 24 Z	2001	121 121 12	177 178 179 1 CAC ACC ATC C D T I	227 22	276 277 278 7 COC COA CAA C C R E
<b>≈</b> 85 ×	26 ×	S 138 138 138 138 138 138 138 138 138 138	55.55	22 SZ ]	222
8 <u>8</u> 5	75 10 10 10 10 10 10 10 10 10 10 10 10 10	75.50 7.00 8.00 8.00 8.00 8.00 8.00 8.00 8.0	1 2	90.00	22
×8~	S S	200-	174 175 1 MCC ATG 1 T M	2 223 224 22 2 TAC 00C A1 Y A Y	273 274 275 7 GCC AAG GAA ( E K E
282	28.38.34	123 124 176 A16 (	25.32 °	223	227
~ <u>6</u> 6 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7	7. 7. 7. 7. N. Y.	25.0	121	727 1 336 1 36	272 Z ATC G
- Se -	~ ¥ ×	283	= 3 ×	280	167
2 <u>C</u> S	2 S u	82>	<u>8</u> 8 ~	219 220 2 000 AMG T A K	270 271 CKG TTC 0 F
- g -	88 — — 88	E 2 -	22 E —	200 V	3.57.788.289.7 3.46.010.010.010 7. L Y
≖໘ີ	28 <u>28</u> ~	≈ <u>;</u> -	88 £ ×	## >	28 H —
≃ 2 >	5 ×	= 3 =	167 168 GM GTG F Y	280	S 32 ×
으 얼 ㅡ	<b>多</b>	5 116 117 118 119 1 ACC ACC CTC C S N V L	節節~	282	第 3 ==
£ 8 ~	28 E ~	===	经第一	1. 212 213 214 215 216 217 218 21 6. 020 602 602 605 606 60 7. P. A. A. S. P. Q. V. J	28 34 >-
±8~	돌은노	<b>≛</b> ‰ ⊸	<u> </u>	7 88 7	医第二
	요 중 ㅡ	SE 35 >	三 完 >	233 A	領領。
W 25 Z	885		162 163 16 1 AM G16 70 1 V	- 8 X	が説め
= 8 -	<del>-</del> 84	AIC CCC // N	들 얼 ~	<b>=3-</b>	至36~
<u>= ≅</u> ∽	880	은 <b>일 =</b>	25 55 N 55 150 5 55 N	1209 210 211 2 000 001 000 0 0 A D	200 261 CTC CTC L V
- E	8350	윤봉스	윤육~	2335	2334
∞ ≩ ≈	∞ = -	観用上	88 第 ~		<b>≋</b> ⊨ –
~ \$ ~	25 N		155 156 157 158 CAC ATA CIG AAG 0 1 L K Peptide 13	206 207 208 2 TCC ATC ATC (	25 F E E
- ¥ <b>=</b>	88 = -	울물 =	35 156 15 0       Peptide	がほく	別屋 ~
~ <b>8 &gt;</b>	8 본 ~	등 등 ~	문용 e e	204 205 2 200 246 2 3 K 1 15 K 1	£ 55 ×
20 ADS GEC ()	조용 -	호유노	\$2 € ==	領別へも	¥2 22 <b>×</b>
~ 5 ~	2 S -	≅ 8 °	E 55 -4	ఐ의용	25 10 5 5
7887	25.55	≅ <u>%</u> ~	2 E -	Per T T Z	Z 3 =
- # =	~ <b>%</b> ~	58°	25 34 ×	232	152 J
. mpoo		uppo	шро	Residue No. Nucleotide codon Anino ocid	поро
로 :활 :등	호 음 등 당	ere. Gide	로 <u>흥</u> 등	로 <u>흥</u> 등	8 15 15 15 15 15 15 15 15 15 15 15 15 15
Residue No. Nucleotide o Animo ocid	Residue No. Nucleotide c Anino ocid		Residue No. Nucleotide c Anino ocid	Residue Mucleolic Anino oci	Residue No. Nucleotide c Anino ocid
중 등 厘	PS 55	Residu Nucleo Anino	窓置道	変置運	Residu Nucleo Anino

FIG.2/

3 1

